


Prior Lines of Therapy Key Requirements for UPLIFT¹



¹Please refer to the protocol Version 9.0 for complete eligibility criteria; Patients in Sweden and France must meet additional criteria for prior lines of therapy

²Please see UPLIFT cohort inclusion criterion #3 for questions on definitions for prior lines

³Based on date of the radiographic image showing progression; if radiographic PD was not documented, date of progression based on biopsy can be used

Key General Requirements for UPLIFT Eligibility¹

- Greater than 18 yo?
- ECOG status 0 or 1?
- Measurable disease by RECIST 1.1?
- Stable treated brain mets?
- Prior toxicities all ≤ grade 1? (except G2 neuropathy)
- LVEF ≥ 50%?
- Adequate organ function?
 - ANC ≥ 1500
 - Platelets > 100,000
 - Hemoglobin > 9.0
 - PT/PTT/INR < 1.2 X ULN
 - GFR ≥ 45
 - Total Bilirubin ≤ ULN
 - AST/ALT ≤ 1.5 ULN
 - Albumin ≥ 3.0
- HBV/HCV negative?

- Room air O2 sat > 93%?
- QTcF < 480 msec?
- Con meds without DILI?
- Con meds without strong CYP450 inhibitors?
- Without liver disease?
- No ongoing severe, uncontrolled systemic disease or intercurrent illness?
- No h/o or active corneal disease?
- No h/o or suspected ILD? No O2 use?
- Last systemic anticancer therapy >28 days prior to planned C1D1
- Prior cancer treated > 2 years prior to screening?
- Archival tissue or new biopsy available?

If any are “no,” then the patient is not eligible

If all are “yes,” then the patient may be eligible for UPLIFT

UPLIFT QTC Substudy

- Con meds without CYP3A4 inducers?
- No uncontrolled cardiac arrhythmias?
- No moderate or severe heart valve disease?
- Sinus rhythm with HR > 45 bpm and < 100 bpm?
- ECG has measurable QT interval?

If any are “No,” then the patient is not eligible for the QTC substudy, but may still be eligible for the general study